



Contents lists available at ScienceDirect

Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis

journal homepage: www.elsevier.com/locate/molmut
Community address: www.elsevier.com/locate/mutres



Review

Extracellular microRNA: A new source of biomarkers

Alton Etheridge, Inyoul Lee, Leroy Hood, David Galas, Kai Wang*

Institute for Systems Biology, 1441 North 34th Street, Seattle, WA 98103, United States

ARTICLE INFO

Article history:

Received 2 December 2010
Received in revised form 3 March 2011
Accepted 7 March 2011
Available online xxx

Keywords:

MicroRNA
Biomarker
Extracellular
Measurement
Body fluid
Export

ABSTRACT

MicroRNAs (miRNAs) are a recently discovered class of small, non-coding RNAs that regulate protein levels post-transcriptionally. miRNAs play important regulatory roles in many cellular processes, including differentiation, neoplastic transformation, and cell replication and regeneration. Because of these regulatory roles, it is not surprising that aberrant miRNA expression has been implicated in several diseases. Recent studies have reported significant levels of miRNAs in serum and other body fluids, raising the possibility that circulating miRNAs could serve as useful clinical biomarkers. Here, we provide a brief overview of miRNA biogenesis and function, the identification and potential roles of circulating extracellular miRNAs, and the prospective uses of miRNAs as clinical biomarkers. Finally, we address several issues associated with the accurate measurement of miRNAs from biological samples.

© 2011 Elsevier B.V. All rights reserved.

Contents

1. miRNA biogenesis and function	00
2. Origin and function of circulating miRNAs	00
3. miRNA potential as biomarkers	00
4. Issues associated with miRNA measurement	00
Conflict of interest	00
Acknowledgements	00
References	00

1. miRNA biogenesis and function

miRNAs are short (19–24 nucleotides in length) non-coding RNAs that regulate messenger RNA (mRNA) or protein levels either by promoting mRNA degradation or by attenuating protein translation. Based on computational prediction, it has been estimated that more than 60% of mammalian mRNAs are targeted by at least one miRNA [1]. miRNAs were first identified in *Caenorhabditis elegans* in the early 1990s [2], but have since been reported in a wide variety of organisms ranging from single-cell algae to humans, suggesting that miRNA-mediated biological function is an ancient and critical cellular regulatory element [3,4]. The importance of miRNA function is further suggested by the extreme evolutionary conservation of both individual miRNA sequences and the miRNA processing machinery. In addition, the number of miRNAs in the

genome appears to be correlated with the complexity of the developmental program, with mammals having the largest number of miRNAs.

miRNAs are transcribed in much the same way as protein-coding genes. The majority of miRNAs are transcribed by RNA polymerase II [5], though a minor fraction of miRNAs that lie within repetitive elements in the genome are transcribed by RNA polymerase III [6]. These primary miRNA transcripts (pri-miRNAs) are often several hundred nucleotides long and are modified similarly to protein-coding transcripts by the addition of a 5' cap and a 3' poly-A tail. Pri-miRNAs are then processed first in the nucleus and later in the cytosol by the RNase III enzymes Drosha and Dicer, respectively. This sequential processing of pri-miRNAs first yields a miRNA precursor (pre-miRNA) of ~70 nucleotides in length, and eventually a mature double-stranded miRNA of 19–24 nucleotides (Fig. 1).

Typically, one strand of this mature miRNA duplex, termed the guide strand, associates with the RNA-induced silencing complex (RISC). While it is generally believed that upon incorporation into the RISC complex, the other strand (the passenger strand)

* Corresponding author.

E-mail address: kwang@systemsbiology.org (K. Wang).




Fig. 1. The biogenesis of miRNAs. miRNA biogenesis begins with transcription of pri-miRNA transcripts by RNA polymerase II or III (1). In the nucleus, pri-miRNAs are processed by Droscha to produce pre-miRNA hairpins (2), which are then exported into the cytosol (3). Here, pre-miRNA hairpins are processed into 19–24 nucleotide mature miRNA duplexes by Dicer (4). One strand of the mature miRNA duplex is incorporated into the RISC complex where it can regulate expression of target mRNAs (5). The other strand may either be degraded, or possibly prepared for export from the cell (6). Some miRNAs have been found packaged in exosomes derived from multivesicular bodies (7). Others may be exported in the presence of RNA-binding proteins (8). Still others might be exported microvesicles shed during membrane blebbing (9). Once in the extracellular space, these miRNAs could be taken up by other cells, degraded by RNases, or excreted (10).

is unwound from the guide strand and degraded, there is evidence that in some cases, both strands of the miRNA duplex are functional [7]. miRNA–RISC complexes interact with mRNA targets through partial sequence complementation, typically within the 3' untranslated region of target mRNAs. It is thought that the extent of base pairing between the miRNA and its mRNA target determines whether the mRNA is degraded or translationally repressed [8].

As with mRNAs, some miRNAs are differentially expressed among tissues or developmental stages. Examples include (i) miR-122, which is preferentially expressed in the liver [9], (ii) miR-133a and miR-133b, which are highly enriched in muscle [10], and (iii) the miR-302 family members which are specific to stem cells [11]. Unlike some widely expressed miRNAs, these tissue- or developmental stage-specific miRNAs likely play key roles in regulating specific processes involved in the development or function of individual tissues. The liver-specific miR-122, for instance, has been implicated in lipid and cholesterol metabolism, which are both known to be important functions of the liver [12]. Because of their restricted expression profiles, these miRNAs hold promise as diagnostic markers or therapeutic targets for tissue- or biological stage-specific diseases. For example, miR-122 is required in hepatitis C virus (HCV) replication [13] and reagents that can modulate the level of miR-122 have moved into clinical development for HCV treatment [13–19].

2. Origin and function of circulating miRNAs

While the majority of miRNAs are found intracellularly, a significant number of miRNAs have been observed outside of cells, including various body fluids [20–24]. These miRNAs are stable

and show distinct expression profiles among different fluid types. Given the instability of most RNA molecules in the extracellular environment, the presence and apparent stability of miRNAs here is surprising. Serum and other body fluids are known to contain ribonucleases [25], which suggests that secreted miRNAs are likely packaged in some manner to protect them against RNase digestion. miRNAs could be shielded from degradation by packaging in lipid vesicles, in complexes with RNA-binding proteins, or both [26,27]. Despite accumulating evidence for the presence of miRNAs in body fluids, the origin and especially the function of these circulating extracellular miRNAs remains poorly understood. One of the more intriguing ideas is that extracellular miRNAs are used as mediators of cell–cell communication [22,28–30]. If this is the case, then certain miRNAs are presumably targeted for export in one cell, and can be recognized, taken up, and utilized by another.

Using cell lines in culture, we have studied the miRNA export phenomenon in detail [22]. Our results revealed distinct intra- and extracellular miRNA spectra, which suggests the existence of a specific miRNA export system. Similar findings have been demonstrated by Pigati et al. in mammary epithelial cells, where the bulk of miR-451 and miR-1246 produced by malignant mammary epithelial cells, but not by non-malignant mammary epithelial cells, was released into the culture medium [31]. In addition to extracellular miRNAs, we have identified several RNA-binding proteins in the medium of cells cultured in the absence of serum [22]. In *in vitro* assays, one of these RNA-binding proteins, nucleophosmin 1 (NPM1), was able to protect synthetic miR-122 miRNA from RNase A digestion. However, further work is needed to determine whether other miRNA-protective proteins exist, as well as whether the apparent NPM1–miR-122 complex is sequence-

Table 1
Characteristics of an ideal biomarker.

Specific
Specific to diseased organ or tissue
Able to differentiate pathologies
Sensitive
Rapid and significant release upon the development of pathology
Predictive
Long half-life in sample
Proportional to degree of severity of pathology
Robust
Rapid, simple, accurate and inexpensive detection
Unconfounded by environment and unrelated conditions
Translatable
Data can be used to bridge pre-clinical and clinical results
Non-invasive
Present in accessible fluid sample

specific or whether NPM1 acts as a general escort for miRNAs. Recently, Kosaka et al. showed that the release of miRNAs is controlled by neutral sphingomyelinase 2 (nSMase2) and through ceramide-dependent secretory machinery [32]. These results suggest the existence of miRNA export system and protein chaperones that could selectively export miRNAs and protect them in the extracellular environment.

Recent studies have also identified both mRNAs and miRNAs in two types of cell-derived lipid vesicles; microvesicles and exosomes. Microvesicles are relatively large (~100 nm to 1 μm) vesicles released from the cell through blebbing. Exosomes, on the other hand, are smaller vesicles (~30–100 nm) released when endosomally derived multivesicular bodies fuse with the plasma membrane. miRNAs have been identified in both exosomes and microvesicles derived from a variety of sources, including human and mouse mast cells [27], glioblastoma tumors [33], plasma [34], saliva [35] and urine [36]. It has recently been reported that miR-150 secreted in microvesicles from human blood cells or cultured THP-1 cells can be taken up by HMEC-1 microvascular endothelial cells and can regulate the expression of c-Myb, a known miR-150 target [37]. Furthermore, Pegtel et al. have shown that miRNAs released in exosomes by Epstein–Barr virus (EBV)-infected cells can be taken up by peripheral blood mononuclear cells and can suppress confirmed EBV target genes [38]. These findings strongly support that at least some exported miRNAs are used for cell-to-cell communication, although much more study is needed to determine how miRNAs are specifically targeted for secretion, recognized for uptake, and what information can be transmitted via this process.

3. miRNA potential as biomarkers

The ideal biomarker should fit a number of criteria depending on how the biomarker is to be used (Table 1). It should be accessible through non-invasive methods, specific to the disease or pathology of interest, a reliable indication of disease before clinical symptoms appear (early detection), sensitive to changes in the pathology (disease progression or therapeutic response), and easily translatable from model systems to humans. Most of the current blood biomarkers are based on the levels of specific proteins in the blood, such as troponin for cardiovascular conditions, carcinoembryonic antigen (CAE) for various cancers, prostate specific antigen (PSA) for prostate cancer, and aminotransferases (alanine aminotransferase, ALT and aspartate aminotransferase, AST) for liver function. Challenges for developing new protein-based biomarkers include the complexity of protein composition in blood, the diversity of post-translational modifications, the low relative abundance of many proteins of interest, the sequence variations among different clinically relevant species, and the difficulties in developing suitable high-affinity detection agents. While proteins are more diverse




Fig. 2. Plasma miR-499 levels in patients with myocardial infarction. The concentration of plasma troponin (bars) and miR-499 (gray squares) was determined from 6 individuals (X-axis) whom may suffer myocardial infarction. The concentration of troponin was displayed as ng/ml and the level of miR-499 was presented in Δ Ct value (40-Ct).

and therefore potentially more informative, the challenges listed above have made the discovery and development of new protein-based biomarkers with proper characteristics an expensive and time-consuming task.

On the other hand, secreted miRNAs have many requisite features of good biomarkers. miRNAs are stable in various bodily fluids, the sequences of most miRNAs are conserved among different species, the expression of some miRNAs is specific to tissues or biological stages, and the level of miRNAs can be easily assessed by various methods, including methods such as polymerase chain reaction (PCR), which allows for signal amplification. The changes of several miRNA levels in plasma, serum, urine, and saliva have already been associated with different diseases [39–59] (Table 2). For example, serum levels of miR-141, have been used to discriminate patients with advanced prostate cancer from healthy individuals [41], the ratio of miR-126 and miR-182 in urine samples can be used to detect bladder cancer [39], and decreased levels of miR-125a and miR-200a in saliva is associated with oral squamous cell carcinoma [50]. In addition to these potential uses in detection of various cancers, another intriguing possibility is the use of levels of organ-specific miRNAs in body fluids to monitor the physiopathological conditions of specific organs.

Using acetaminophen overdose as a model, we have previously demonstrated the possibility of using specific miRNA levels in blood to detect drug-induced liver injury [60]. The severity of liver injury in this well-established model system can be precisely detected and monitored by measuring the levels of miR-122, a liver-specific miRNA, in plasma using quantitative PCR (qPCR). While elevated plasma ALT levels usually indicate impairment of liver function, plasma ALT originating from extra-hepatic sources have also been reported from patients with burns, muscle inflammation, hypothyroidism, or myopathies [61]. This makes the use of ALT levels alone in diagnosis of liver diseases unreliable, whereas the expression of miR-122 is fairly restricted to liver. In our hands, this miRNA-based method is more sensitive and probably more reliable than the current serological method, plasma ALT level, to detect and monitor drug-induced liver injuries. Besides miR-122 for liver injury, we and others [62] have demonstrated that the plasma level of miR-499, a heart specific miRNA, shows a perfect correlation with blood troponin levels in patients with myocardial infarction (Fig. 2). This clearly demonstrates the possibility of using the level of a tissue-specific miRNA in circulation to reflect the health status of targeted tissues.

4. Issues associated with miRNA measurement

As mentioned above, extracellular circulating miRNAs offer several potential advantages as informative biomarkers when

Table 2
Circulating microRNAs proposed as diagnostic biomarkers in human diseases.

Type of cancer	Biomarker candidate	References
Prostate cancer	Fifteen serum miRNAs were over-expressed from stage 3 and 4 prostate cancer patients (miR-16, -92a, -103, -107, -197, -34b, -328, -485-3p, -486-5p, -92b, -574-3p, -636, -640, -766, -885-5p) when compared to normal individuals	[40]
	Expression levels of <i>miR-141</i> in serum can distinguish prostate cancer patients from healthy controls	[41]
	Forty-eight serum miRNAs were differentially expressed in breast cancer patients (22 up-regulated, 26 down-regulated) when compared to controls	[42]
Breast cancer	Increased expression levels of <i>miR-10b</i> and <i>-34a</i> in serum was observed in breast cancer patients	[43]
	Decreased expression levels of <i>miR-195</i> and <i>let-7a</i> in serum was observed in breast cancer patients	[44]
Ovarian cancer	<i>miR-21</i> , <i>92</i> , <i>93</i> , <i>126</i> and <i>29a</i> were over-expressed in serum samples from cancer patients compared to controls and <i>miR-155</i> , <i>127</i> and <i>99b</i> were under-expressed	[52]
	Eight serum exosomal miRNAs were elevated in ovarian cancer patients: <i>miR-21</i> , <i>-141</i> , <i>-200a</i> , <i>-200b</i> , <i>-200c</i> , <i>-203</i> , <i>-205</i> , <i>-214</i> .	[45]
Lung cancer	Eleven serum miRNAs (including <i>miR-7i</i> , <i>-146b</i> , <i>-206</i> , and <i>-21</i>) were changed more than five-fold by NGS between longer-survival lung cancer patient groups and shorter-survival groups. Levels of four miRNAs (<i>miR-486</i> , <i>-30d</i> , <i>-1</i> , <i>-499</i>) were associated with overall survival	[46]
Colorectal cancer	Both <i>miR-29a</i> and <i>miR-92a</i> showed elevated levels in plasma from 37 advanced colorectal cancer patients.	[47]
	The levels of two plasma miRNAs (<i>miR-17-3p</i> and <i>-92a</i>) were significantly elevated.	[51]
Bladder cancer	The ratio of two urinary miRNAs (<i>miR-126</i> and <i>-182</i>) enabled detection of urinary bladder cancer.	[39]
Oral cancer	The level of plasma <i>miR-31</i> was significantly elevated in oral squamous cell carcinoma patients compared to the control groups.	[72]
	Two miRNAs (<i>miR-125a</i> and <i>-200a</i>) showed decreased levels in saliva with oral cancer	[50]
Heart conditions	From acute myocardial infarction patients, the level of plasma <i>miR-208b</i> and <i>-499</i> was highly elevated and correlated with plasma troponin level.	[53]
	Level of six plasma miRNAs including <i>miR-423-5p</i> was elevated in patients with heart failure.	[54]
	Plasma <i>miR-1</i> level was significantly elevated from patients with acute myocardial infarction.	[55,56]

compared to protein-based blood biomarkers. For instance, while low abundance can significantly hinder the detection of some protein-based biomarkers, most circulating miRNAs can be readily detected by PCR. Additionally, protein-based biomarkers may have different post-translational modifications which can affect the accuracy of measurement, but miRNA species are relatively homogenous. Furthermore, the highly specific expression profile of several miRNAs such as miR-122 in the liver and miR-499 in

Table 3
Issues associated with miRNA measurement.

Measurement
Low correlation between different measurement platforms
Short and conserved sequences in paralogs make it difficult to measure specific miRNA levels
Difficult to distinguish between precursor and mature forms
Sample
Lack of standard protocols for sample preparation
Concentration measurement for miRNA in sample is difficult
Data processing
Normalization among different samples, especially for extracellular miRNA
Difficult to validate results especially when using different protocols and platforms

the heart add the possibility of using the levels of these miRNAs in blood to precisely monitor the health status of specific organs. However, to further develop miRNA-based biomarkers, there are several fundamental issues associated with miRNA measurements that still need to be addressed (Table 3).

As with mRNA measurement, there are two major global miRNA profiling platforms in use today; microarrays and qPCR-based methods. Each of these methods has benefits and drawbacks. Although qPCR methods for miRNA quantitation are relatively inexpensive, widely available, and allow measurements of very small quantities of miRNAs, the primer design can heavily influence the results. Microarray-based measurement methods generally require more starting material than qPCR, and it can be challenging to develop probes and hybridization conditions that work well to detect many different miRNAs at once. In addition to microarrays and qPCR, there are other less frequently used methods, such as traditional northern blotting, ligation based measurement [63], and direct sequencing using next generation sequencing (NGS) platforms [64]. Of these methods, miRNA profiling by NGS may be the most promising, as it largely avoids many miRNA measurement pitfalls. The use of NGS technology offers the possibility to obtain comprehensive and accurate measurement for transcripts as well as miRNAs. Since the NGS approach is largely sequence independent, it does not rely on the design of primers or probes specific to each miRNA. Instead, NGS sequencing provides the number of counts for each miRNA or transcript present in the sample. Despite these advantages, NGS remains expensive and labor intensive, both in the sample preparation and in data analysis.

For miRNA measurement, it is well acknowledged that there is low correlation of results obtained from different platforms or even from the same platform using products from different vendors (such as microarrays from Agilent and Affymetrix) [65,66]. The short length and high sequence similarity among some miRNAs likely contributes to the inconsistency of measurement results due to the problems in designing specific primers for qPCR or probes for microarrays. Additionally, sequencing of miRNAs has revealed significant sequence heterogeneity at the 3' and 5' ends (termed isomirs) that may further complicate measurements [67]. This miRNA sequence variation probably results from imprecise processing of miRNAs. However, our preliminary results indicate that there may be functional association with differences in the abundance of individual isomirs [67]. Further supporting the possibility of functionally significant isomirs, dynamic changes in the isomir population have been observed during *Drosophila* development [68]. Using synthetic isomir sequences, we have shown that sequence heterogeneity at the ends of isomirs may further complicate the miRNA measurement results, especially for some qPCR-based methods [67]. We have created an isomir database (<http://galas.systemsbiology.net/cgi-bin/isomir/find.pl>), to collect observed miRNA sequence variations which we hope may facilitate better miRNA probe or primer design in the future.

Another critical issue associated with miRNA measurement is assessing the quantity and quality of miRNA isolated from biological samples. In contrast to isolated mRNA, where the sizes and relative abundance of ribosomal RNAs can be used to assess the integrity of the RNA preparation, there is currently no good way to gauge the quality of isolated miRNA. In addition, degraded mRNA can interfere with the quantitation of miRNA preparations. Measuring the amount of miRNA isolated from body fluid samples is especially difficult due to the low concentration of RNA typically obtained from these samples. Therefore, it is difficult to establish the amount of input RNA in any measurement platform, which may make sample-to-sample and cross platform comparisons challenging.

Most protein-based blood biomarkers are measured using enzyme linked immunosorbent assays (ELISA) on serum. EDTA-plasma, on the other hand, is the preferred sample type for blood proteome studies, since it offers more reproducible protein measurement [46,69]. However, for blood miRNA measurements, both plasma and serum have been used in various studies (examples see Table 2). Our preliminary results suggest that there are notable differences in miRNA levels between serum and plasma. However, a comprehensive and systematic comparison is needed to establish the difference, if any, between serum and plasma miRNA content, as well as the effects of various anticoagulants on miRNA measurement.

Another issue that remains to be addressed is how to normalize or compare miRNA measurement results between samples. There are several well-established protocols for normalization of gene expression measurements across different samples. These include using global normalization approaches such as the level of housekeeping genes. Similar approaches can be used to normalize cellular miRNA measurements. However, for extracellular miRNAs, there are currently no known extracellular housekeeping RNAs that can be used for normalization. In many cases, normalization by input volume remains the best option when measuring circulating miRNAs. One interesting possibility is to use the level of other biomolecules in body fluids such as creatinine levels in blood and urine to normalize miRNAs. The levels of creatinine have been used to “normalize” other biomolecules in urine, and it provides an interesting possibility as a normalization basis for miRNAs in plasma and urine [70,71].

Finding informative biomarkers is not only key to understanding physiopathological processes of diseases, but is also critical for therapeutic development. It has been one of the major focuses of biomedical research in the past two decades. Despite the identification of several new protein-based biomarkers, very few of them actually pass rigorous validation processes. Circulating extracellular miRNAs clearly possess many desirable properties when compared to protein biomarkers. However, they require the same stringent validation processes as protein-based biomarkers to demonstrate their specificity and selectivity. The most fundamental challenges in miRNA-based biomarker development are the issues associated with accurate measurement. Building better reagent sets is one of the key steps to obtaining more comprehensive and accurate miRNA measurements. Setting up a standardization process for sample preparation and developing a more accurate method to assess the quality and quantity of miRNA are also urgently needed.

Conflict of interest

None.

Acknowledgements

This work was supported by the ISB-University of Luxembourg program, a grant from the National Prion Research Program

(NS41997), research subcontract from Georgetown University (RX 4222-802-ISB) and research contracts from the Department of Defense (W911SR-07-C-0101, W911NF-09-D-0001 and HDTRA 1-08-C-0023).

References

- [1] R.C. Friedman, K.K. Farh, C.B. Burge, D.P. Bartel, Most mammalian mRNAs are conserved targets of microRNAs, *Genome Res.* 19 (2009) 92–105.
- [2] R.C. Lee, R.L. Feinbaum, V. Ambros, The *C. elegans* heterochronic gene *lin-4* encodes small RNAs with antisense complementarity to *lin-14*, *Cell* 75 (1993) 843–854.
- [3] D.P. Bartel, C.Z. Chen, Micromanagers of gene expression: the potentially widespread influence of metazoan microRNAs, *Nat. Rev. Genet.* 5 (2004) 396–400.
- [4] T. Zhao, G. Li, S. Mi, S. Li, G.J. Hannon, X.J. Wang, Y. Qi, A complex system of small RNAs in the unicellular green alga *Chlamydomonas reinhardtii*, *Genes Dev.* 21 (2007) 1190–1203.
- [5] Y. Lee, M. Kim, J. Han, K.H. Yeom, S. Lee, S.H. Baek, V.N. Kim, MicroRNA genes are transcribed by RNA polymerase II, *EMBO J.* 23 (2004) 4051–4060.
- [6] G.M. Borchert, W. Lanier, B.L. Davidson, RNA polymerase III transcribes human microRNAs, *Nat. Struct. Mol. Biol.* 13 (2006) 1097–1101.
- [7] H.Y. Hu, Z. Yan, Y. Xu, H. Hu, C. Menzel, Y.H. Zhou, W. Chen, P. Khaitovich, Sequence features associated with microRNA strand selection in humans and flies, *BMC Genomics* 10 (2009) 413.
- [8] M.R. Fabian, N. Sonenberg, W. Filipowicz, Regulation of mRNA translation and stability by microRNAs, *Annu. Rev. Biochem.* 79 (2010) 351–379.
- [9] M. Lagos-Quintana, R. Rauhut, A. Yalcin, J. Meyer, W. Lendeckel, T. Tuschl, Identification of tissue-specific microRNAs from mouse, *Curr. Biol.* 12 (2002) 735–739.
- [10] L.F. Sempere, S. Freemantle, I. Pitha-Rowe, E. Moss, E. Dmitrovsky, V. Ambros, Expression profiling of mammalian microRNAs uncovers a subset of brain-expressed microRNAs with possible roles in murine and human neuronal differentiation, *Genome Biol.* 5 (2004) R13.
- [11] A. Barroso-del Jesus, G. Lucena-Aguilar, P. Menendez, The miR-302-367 cluster as a potential stemness regulator in ESCs, *Cell Cycle* 8 (2009) 394–398.
- [12] C. Esau, S. Davis, S.F. Murray, X.X. Yu, S.K. Pandey, M. Pear, L. Watts, S.L. Booten, M. Graham, R. McKay, A. Subramaniam, S. Propp, B.A. Lollo, S. Freier, C.F. Bennett, S. Bhanot, B.P. Monia, miR-122 regulation of lipid metabolism revealed by in vivo antisense targeting, *Cell Metab.* 3 (2006) 87–98.
- [13] C.L. Jopling, M. Yi, A.M. Lancaster, S.M. Lemon, P. Sarnow, Modulation of hepatitis C virus RNA abundance by a liver-specific microRNA, *Science* 309 (2005) 1577–1581.
- [14] S. Pfeffer, T.F. Baumert, Impact of microRNAs for pathogenesis and treatment of hepatitis C virus infection, *Gastroenterol. Clin. Biol.* 34 (2010) 431–435.
- [15] D.D. Young, C.M. Connelly, C. Grohmann, A. Deiters, Small molecule modifiers of microRNA miR-122 function for the treatment of hepatitis C virus infection and hepatocellular carcinoma, *J. Am. Chem. Soc.* 132 (2010) 7976–7981.
- [16] R.E. Lanford, E.S. Hildebrandt-Eriksen, A. Petri, R. Persson, M. Lindov, M.E. Munk, S. Kauppinen, H. Orum, Therapeutic silencing of microRNA-122 in primates with chronic hepatitis C virus infection, *Science* 327 (2010) 198–201.
- [17] M. Niepmann, Activation of hepatitis C virus translation by a liver-specific microRNA, *Cell Cycle* 8 (2009) 1473–1477.
- [18] J. Lupberger, L. Brino, T.F. Baumert, RNAi: a powerful tool to unravel hepatitis C virus-host interactions within the infectious life cycle, *J. Hepatol.* 48 (2008) 523–525.
- [19] Q.W. Pan, S.D. Henry, B.J. Scholte, H.W. Tilanus, H.L. Janssen, L.J. van der Laan, New therapeutic opportunities for hepatitis C based on small RNA, *World J. Gastroenterol.* 13 (2007) 4431–4436.
- [20] K. Zen, C.Y. Zhang, Circulating MicroRNAs: a novel class of biomarkers to diagnose and monitor human cancers, *Med. Res. Rev.* (2010).
- [21] J.A. Weber, D.H. Baxter, S. Zhang, D.Y. Huang, K.H. Huang, M.J. Lee, D.J. Galas, K. Wang, The microRNA spectrum in 12 body fluids, *Clin. Chem.* 56 (2010) 1733–1741.
- [22] K. Wang, S. Zhang, J. Weber, D. Baxter, D.J. Galas, Export of microRNAs and microRNA-protective protein by mammalian cells, *Nucleic Acids Res.* 38 (2010) 7248–7259.
- [23] D. Zubakov, A.W. Boersma, Y. Choi, P.F. van Kuijk, E.A. Wiemer, M. Kayser, MicroRNA markers for forensic body fluid identification obtained from microarray screening and quantitative RT-PCR confirmation, *Int. J. Legal Med.* 124 (2010) 217–226.
- [24] E.K. Hanson, H. Lubenow, J. Ballantyne, Identification of forensically relevant body fluids using a panel of differentially expressed microRNAs, *Anal. Biochem.* 387 (2009) 303–314.
- [25] J.L. Weickmann, D.G. Glitz, Human ribonucleases. Quantitation of pancreatic-like enzymes in serum, urine, and organ preparations, *J. Biol. Chem.* 257 (1982) 8705–8710.
- [26] D.J. Gibbins, C. Ciaudo, M. Erhardt, O. Voinnet, Multivesicular bodies associate with components of miRNA effector complexes and modulate miRNA activity, *Nat. Cell Biol.* 11 (2009) 1143–1149.

- [27] H. Valadi, K. Ekstrom, A. Bossios, M. Sjostrand, J.J. Lee, J.O. Lotvall, Exosome-mediated transfer of mRNAs and microRNAs is a novel mechanism of genetic exchange between cells, *Nat. Cell Biol.* 9 (2007) 654–659.
- [28] H. Iguchi, N. Kosaka, T. Ochiya, Secretory microRNAs as a versatile communication tool, *Commun. Integr. Biol.* 3 (2010) 478–481.
- [29] G. Camussi, M.C. Deregiibus, S. Bruno, V. Cantaluppi, L. Biancone, Exosomes/microvesicles as a mechanism of cell-to-cell communication, *Kidney Int.* 78 (2010) 838–848.
- [30] V. Muralidharan-Chari, J.W. Clancy, A. Sedgwick, C. D'Souza-Schorey, Microvesicles: mediators of extracellular communication during cancer progression, *J. Cell Sci.* 123 (2010) 1603–1611.
- [31] L. Pigati, S.C. Yaddanapudi, R. Iyengar, D.J. Kim, S.A. Hearn, D. Danforth, M.L. Hastings, D.M. Duelli, Selective release of microRNA species from normal and malignant mammary epithelial cells, *PLoS One* 5 (2010) e13515.
- [32] N. Kosaka, H. Iguchi, Y. Yoshioka, F. Takeshita, Y. Matsuki, T. Ochiya, Secretory mechanisms and intercellular transfer of microRNAs in living cells, *J. Biol. Chem.* 285 (2010) 17442–17452.
- [33] J. Skog, T. Wurdinger, S. van Rijn, D.H. Meijer, L. Gainche, M. Sena-Esteves, W.T. Curry Jr., B.S. Carter, A.M. Krichevsky, X.O. Breakefield, Glioblastoma microvesicles transport RNA and proteins that promote tumour growth and provide diagnostic biomarkers, *Nat. Cell Biol.* 10 (2008) 1470–1476.
- [34] M.P. Hunter, N. Ismail, X. Zhang, B.D. Aguda, E.J. Lee, L. Yu, T. Xiao, J. Schafer, M.L. Lee, T.D. Schmittgen, S.P. Nana-Sinkam, D. Jarjoura, C.B. Marsh, Detection of microRNA expression in human peripheral blood microvesicles, *PLoS One* 3 (2008) e3694.
- [35] A. Michael, S.D. Bajracharya, P.S. Yuen, H. Zhou, R.A. Star, G.G. Illei, I. Alevizos, Exosomes from human saliva as a source of microRNA biomarkers, *Oral Dis.* 16 (2010) 34–38.
- [36] I. Dimov, L. Jankovic Velickovic, V. Stefanovic, Urinary exosomes, *ScientificWorldJournal* 9 (2009) 1107–1118.
- [37] Y. Zhang, D. Liu, X. Chen, J. Li, L. Li, Z. Bian, F. Sun, J. Lu, Y. Yin, X. Cai, Secreted monocytic miR-150 enhances targeted endothelial cell migration, *Mol. Cell* 39 (2010) 133–144.
- [38] D.M. Pegtel, K. Cosmopoulos, D.A. Thorley-Lawson, M.A. van Eijndhoven, E.S. Hopmans, J.L. Lindenberg, T.D. de Gruijil, T. Wurdinger, J.M. Middeldorp, Functional delivery of viral miRNAs via exosomes, *Proc. Natl. Acad. Sci. U. S. A.* 107 (2010) 6328–6333.
- [39] M. Hanke, K. Hoefig, H. Merz, A.C. Feller, I. Kausch, D. Jocham, J.M. Warnecke, G. Szakiel, A robust methodology to study urine microRNA as tumor marker: microRNA-126 and microRNA-182 are related to urinary bladder cancer, *Urol. Oncol.* 28 (2010) 655–661.
- [40] M.J. Lodes, M. Caraballo, D. Suci, S. Munro, A. Kumar, B. Anderson, Detection of cancer with serum miRNAs on an oligonucleotide microarray, *PLoS One* 4 (2009) e6229.
- [41] P.S. Mitchell, R.K. Parkin, E.M. Kroh, B.R. Fritz, S.K. Wyman, E.L. Pogosova-Agadjanian, A. Peterson, J. Noteboom, K.C. O'Brian, A. Allen, D.W. Lin, N. Urban, C.W. Drescher, B.S. Knudsen, D.L. Stirewalt, R. Gentleman, R.L. Vessella, P.S. Nelson, D.B. Martin, M. Tewari, Circulating microRNAs as stable blood-based markers for cancer detection, *Proc. Natl. Acad. Sci. U.S.A.* 105 (2008) 10513–10518.
- [42] H. Zhao, J. Shen, L. Medico, D. Wang, C.B. Ambrosone, S. Liu, A pilot study of circulating miRNAs as potential biomarkers of early stage breast cancer, *PLoS One* 5 (2010) e13735.
- [43] C. Roth, B. Rack, V. Muller, W. Janni, K. Pantel, H. Schwarzenbach, Circulating microRNAs as blood-based markers for patients with primary and metastatic breast cancer, *Breast Cancer Res.* 12 (2010) R90.
- [44] H.M. Heneghan, N. Miller, A.J. Lowery, K.J. Sweeney, J. Newell, M.J. Kerin, Circulating microRNAs as novel minimally invasive biomarkers for breast cancer, *Ann. Surg.* 251 (2010) 499–505.
- [45] D.D. Taylor, C. Cercel-Taylor, MicroRNA signatures of tumor-derived exosomes as diagnostic biomarkers of ovarian cancer, *Gynecol. Oncol.* 110 (2008) 13–21.
- [46] A.J. Rai, C.A. Gelfand, B.C. Haywood, D.J. Warunek, J. Yi, M.D. Schuchard, R.J. Mehig, S.L. Cockrill, G.B. Scott, H. Tammen, P. Schulz-Knappe, D.W. Speicher, F. Vitthum, B.B. Haab, G. Siest, D.W. Chan, HUPO Plasma proteome project specimen collection and handling: towards the standardization of parameters for plasma proteome samples, *Proteomics* 5 (2005) 3262–3277.
- [47] Z. Huang, D. Huang, S. Ni, Z. Peng, W. Sheng, X. Du, Plasma microRNAs are promising novel biomarkers for early detection of colorectal cancer, *Int. J. Cancer* 127 (2010) 118–126.
- [48] E.K. Ng, W.P. Tsang, S.S. Ng, H.C. Jin, J. Yu, J.J. Li, C. Rocken, M.P. Ebert, T.T. Kwok, J.J. Sung, MicroRNA-143 targets DNA methyltransferases 3A in colorectal cancer, *Br. J. Cancer* 101 (2009) 699–706.
- [49] C.J. Liu, S.Y. Kao, H.F. Tu, M.M. Tsai, K.W. Chang, S.C. Lin, Increase of microRNA miR-31 level in plasma could be a potential marker of oral cancer, *Oral Dis.* 16 (2010) 360–364.
- [50] N.J. Park, H. Zhou, D. Elashoff, B.S. Henson, D.A. Kastratovic, E. Abemayor, D.T. Wong, Salivary microRNA: discovery, characterization, and clinical utility for oral cancer detection, *Clin. Cancer Res.* 15 (2009) 5473–5477.
- [51] E.K. Ng, W.W. Chong, H. Jin, E.K. Lam, V.Y. Shin, J. Yu, T.C. Poon, S.S. Ng, J.J. Sung, Differential expression of microRNAs in plasma of patients with colorectal cancer: a potential marker for colorectal cancer screening, *Gut* 58 (2009) 1375–1381.
- [52] K.E. Resnick, H. Alder, J.P. Hagan, D.L. Richardson, C.M. Croce, D.E. Cohn, The detection of differentially expressed microRNAs from the serum of ovarian cancer patients using a novel real-time PCR platform, *Gynecol. Oncol.* 112 (2009) 55–59.
- [53] M.F. Corsten, R. Dennert, S. Jochems, T. Kuznetsova, Y. Devaux, L. Hofstra, D.R. Wagner, J.A. Staessen, S. Heymans, B. Schroen, Circulating MicroRNA-208b and MicroRNA-499 reflect myocardial damage in cardiovascular disease, *Circ. Cardiovasc. Genet.* 3 (2010) 499–506.
- [54] A.J. Tijssen, E.E. Creemers, P.D. Moerland, L.J. de Windt, A.C. van der Wal, W.E. Kok, Y.M. Pinto, MiR423-5p as a circulating biomarker for heart failure, *Circ. Res.* 106 (2010) 1035–1039.
- [55] J. Ai, R. Zhang, Y. Li, J. Pu, Y. Lu, J. Jiao, K. Li, B. Yu, Z. Li, R. Wang, L. Wang, Q. Li, N. Wang, H. Shan, Z. Li, B. Yang, Circulating microRNA-1 as a potential novel biomarker for acute myocardial infarction, *Biochem. Biophys. Res. Commun.* 391 (2010) 73–77.
- [56] Y. Cheng, N. Tan, J. Yang, X. Liu, X. Cao, P. He, X. Dong, S. Qin, C. Zhang, A translational study of circulating cell-free microRNA-1 in acute myocardial infarction, *Clin. Sci. (Lond.)* 119 (2010) 87–95.
- [57] M. Ferracin, A. Veronese, M. Negrini, Micromarkers: miRNAs in cancer diagnosis and prognosis, *Expert Rev. Mol. Diagn.* 10 (2010) 297–308.
- [58] Y. Tie, B. Liu, H. Fu, X. Zheng, Circulating miRNA and cancer diagnosis, *Sci. China C Life Sci.* 52 (2009) 1117–1122.
- [59] M. Fabbri, miRNAs as molecular biomarkers of cancer, *Expert Rev. Mol. Diagn.* 10 (2010) 435–444.
- [60] K. Wang, S. Zhang, B. Marzolf, P. Troisch, A. Brightman, Z. Hu, L.E. Hood, D.J. Galas, Circulating microRNAs, potential biomarkers for drug-induced liver injury, *Proc. Natl. Acad. Sci. U.S.A.* 106 (2009) 4402–4407.
- [61] M. Mazer, J. Perrone, Acetaminophen-induced nephrotoxicity: pathophysiology, clinical manifestations, and management, *J. Med. Toxicol.* 4 (2008) 2–6.
- [62] T. Adachi, M. Nakanishi, Y. Otsuka, K. Nishimura, G. Hirokawa, Y. Goto, H. Nonogi, N. Iwai, Plasma microRNA 499 as a biomarker of acute myocardial infarction, *Clin. Chem.* 56 (2010) 1183–1185.
- [63] J. Li, B. Yao, H. Huang, Z. Wang, C. Sun, Y. Fan, Q. Chang, S. Li, X. Wang, J. Xi, Real-time polymerase chain reaction microRNA detection based on enzymatic stem-loop probes ligation, *Anal. Chem.* 81 (2009) 5446–5451.
- [64] M.F. Thomas, K.M. Ansel, Construction of small RNA cDNA libraries for deep sequencing, *Methods Mol. Biol.* 667 (2010) 93–111.
- [65] F. Sato, S. Tsuchiya, K. Terasawa, G. Tsujimoto, Intra-platform repeatability and inter-platform comparability of microRNA microarray technology, *PLoS ONE* 4 (2009) e5540.
- [66] Y. Chen, J.A. Gelfond, L.M. McManus, P.K. Shireman, Reproducibility of quantitative RT-PCR array in miRNA expression profiling and comparison with microarray analysis, *BMC Genomics* 10 (2009) 407.
- [67] L.W. Lee, S. Zhang, A. Etheridge, L. Ma, D. Martin, D. Galas, K. Wang, Complexity of the microRNA repertoire revealed by next-generation sequencing, *RNA* 16 (2010) 2170–2180.
- [68] S.L. Fernandez-Valverde, R.J. Taft, J.S. Mattick, Dynamic isomiR regulation in *Drosophila* development, *RNA* 16 (2010) 1881–1888.
- [69] G.S. Omenn, Exploring the human plasma proteome, *Proteomics* 5 (2005) 3223–3225.
- [70] S.L. Goldstein, Urinary kidney injury biomarkers and urine creatinine normalization: a false premise or not? *Kidney Int.* 78 (2010) 433–435.
- [71] B.D. Wagner, F.J. Accurso, T.A. Laguna, The applicability of urinary creatinine as a method of specimen normalization in the cystic fibrosis population, *J. Cyst. Fibros.* 9 (2010) 212–216.
- [72] B. Czech, R. Zhou, Y. Erlich, J. Brennecke, R. Binari, C. Villalta, A. Gordon, N. Perrimon, G.J. Hannon, Hierarchical rules for argonaute loading in *Drosophila*, *Mol. Cell* 36 (2009) 445–456.