

47. Yu N, Christiaens O, Liu J, Niu J, Cappelle K, Caccia S, et al. Delivery of dsRNA for RNAi in insects: an overview and future directions. *Insect Sci* 2012; 20:4-14; <http://dx.doi.org/10.1111/j.1744-7917.2012.01534.x>
48. Amdam GV, Simões ZL, Guidugli KR, Norberg K, Omholt SW. Disruption of vitellogenin gene function in adult honeybees by intra-abdominal injection of double-stranded RNA. *BMC Biotechnol* 2003; 3:1; PMID:12546706; <http://dx.doi.org/10.1186/1472-6750-3-1>
49. Aronstein K, Pankiwi T, Saldivar E. SID-I is implicated in systemic gene silencing in the honey bee. *J Apic Res* 2006; 45:20-4
50. Maori E, Paldi N, Shafir S, Kalev H, Tsur E, Glick E, et al. IAPV, a bee-affecting virus associated with Colony Collapse Disorder can be silenced by dsRNA ingestion. *Insect Mol Biol* 2009; 18:55-60; PMID:19196347; <http://dx.doi.org/10.1111/j.1365-2583.2009.00847.x>
51. Nunes FM, Simões ZL. A non-invasive method for silencing gene transcription in honeybees maintained under natural conditions. *Insect Biochem Mol Biol* 2009; 39:157-60; PMID:19049870; <http://dx.doi.org/10.1016/j.ibmb.2008.10.011>
52. Hunter W, Ellis J, Vanengelsdorp D, Hayes J, Westervelt D, Glick E, et al. Large-scale field application of RNAi technology reducing Israeli acute paralysis virus disease in honey bees (*Apis mellifera*, Hymenoptera: Apidae). *PLoS Pathog* 2010; 6:e1001160; PMID:21203478; <http://dx.doi.org/10.1371/journal.ppat.1001160>
53. Liu X, Zhang Y, Yan X, Han R. Prevention of Chinese sacbrood virus infection in *Apis cerana* using RNA interference. *Curr Microbiol* 2010; 61:422-8; PMID:20379718; <http://dx.doi.org/10.1007/s00284-010-9633-2>
54. Desai SD, Eu YJ, Whyard S, Currie RW. Reduction in deformed wing virus infection in larval and adult honey bees (*Apis mellifera* L.) by double-stranded RNA ingestion. *Insect Mol Biol* 2012; 21:446-55; PMID:22690671; <http://dx.doi.org/10.1111/j.1365-2583.2012.01150.x>
55. Nazzi F, Brown SP, Annoscia D, Del Piccolo F, Di Prisco G, Varricchio P, et al. Synergistic parasite-pathogen interactions mediated by host immunity can drive the collapse of honeybee colonies. *PLoS Pathog* 2012; 8:e1002735; PMID:22719246; <http://dx.doi.org/10.1371/journal.ppat.1002735>
56. Garbian Y, Maori E, Kalev H, Shafir S, Sela I. Bidirectional transfer of RNAi between honey bee and *Varroa* destructor: *Varroa* gene silencing reduces *Varroa* population. *PLoS Pathog* 2012; 8:e1003035; PMID:23308063; <http://dx.doi.org/10.1371/journal.ppat.1003035>
57. Czech B, Hannon GJ. Small RNA sorting: matchmaking for Argonautes. *Nat Rev Genet* 2011; 12:19-31; PMID:21116305; <http://dx.doi.org/10.1038/nrg2916>
58. Crailsheim K, Schneider L, Hrassnigg N, Buhlmann G, Brosch U, Gmeinbauer R, et al. Pollen consumption and utilization in worker honeybees (*Apis mellifera* carnica): dependence on individual age and function. *J Insect Physiol* 1992; 38:409-19; [http://dx.doi.org/10.1016/0022-1910\(92\)90117-V](http://dx.doi.org/10.1016/0022-1910(92)90117-V)
59. Rortais A, Arnold G, Halm MP, Touffet-Briens F. Modes of honeybees exposure to systemic insecticides: estimated amounts of contaminated pollen and nectar consumed by different categories of bees. *Apidologie (Celle)* 2005; 36:71-83; <http://dx.doi.org/10.1051/apido:2004071>
60. Shi YY, Wu XB, Huang ZY, Wang ZL, Yan WY, Zeng ZJ. Epigenetic modification of gene expression in honey bees by heterospecific gland secretions. *PLoS One* 2012; 7:e43727; PMID:22928024; <http://dx.doi.org/10.1371/journal.pone.0043727>
61. Fabbri M, Paone A, Calore F, Galli R, Gaudio E, Santhanam R, et al. MicroRNAs bind to Toll-like receptors to induce prometastatic inflammatory response. *Proc Natl Acad Sci USA* 2012; 109:E2110-6; PMID:22753494; <http://dx.doi.org/10.1073/pnas.1209414109>
62. van Nood E, Vrieze A, Nieuwdorp M, Fuentes S, Zoetendal EG, de Vos WM, et al. Duodenal infusion of donor feces for recurrent *Clostridium difficile*. *N Engl J Med* 2013; 368:407-15; PMID:23323867; <http://dx.doi.org/10.1056/NEJMoa1205037>
63. Lu TX, Hartner J, Lim EJ, Fabry V, Mingler MK, Cole ET, et al. MicroRNA-21 limits in vivo immune response-mediated activation of the IL-12/IFN-gamma pathway, Th1 polarization, and the severity of delayed-type hypersensitivity. *J Immunol* 2011; 187:3362-73; PMID:21849676; <http://dx.doi.org/10.4049/jimmunol.1101235>
64. Chan SY, Zhang YY, Hemann C, Mahoney CE, Zweier JL, Loscalzo J. MicroRNA-210 controls mitochondrial metabolism during hypoxia by repressing the iron-sulfur cluster assembly proteins ISCU1/2. *Cell Metab* 2009; 10:273-84; PMID:19808020; <http://dx.doi.org/10.1016/j.cmet.2009.08.015>
65. Ben-Shahar Y, Robichon A, Sokolowski MB, Robinson GE. Influence of gene action across different time scales on behavior. *Science* 2002; 296:741-4; PMID:11976457; <http://dx.doi.org/10.1126/science.1069911>
66. Schmittgen TD, Lee EJ, Jiang J, Sarkar A, Yang L, Elton TS, et al. Real-time PCR quantification of precursor and mature microRNA. *Methods* 2008; 44:31-8; PMID:18158130; <http://dx.doi.org/10.1016/j.ymeth.2007.09.006>