

siRNA and miRNA: an insight into RISCs

Guiliang Tang

Department of Biochemistry and Molecular Pharmacology, University of Massachusetts Medical School, Worcester, MA 01605, USA

Two classes of short RNA molecule, small interfering RNA (siRNA) and microRNA (miRNA), have been identified as sequence-specific posttranscriptional regulators of gene expression. siRNA and miRNA are incorporated into related RNA-induced silencing complexes (RISCs), termed siRISC and miRISC, respectively. The current model argues that siRISC and miRISC are functionally interchangeable and target specific mRNAs for cleavage or translational repression, depending on the extent of sequence complementarity between the small RNA and its target. Emerging evidence indicates, however, that siRISC and miRISC are distinct complexes that regulate mRNA stability and translation. The assembly of RISCs can be traced from the biogenesis of the small RNA molecules and the recruitment of these RNAs by the RISC loading complex (RLC) to the transition of the RLC into the active RISC. Target recognition by the RISC can then take place through different interacting modes.

Introduction

Although small interfering RNA (siRNA; see Glossary) and microRNA (miRNA) were initially discovered in unrelated studies, both types of small RNA are closely related in their biogenesis, assembly into RNA–protein complexes and ability to regulate gene transcripts negatively in diverse eukaryotes [1–4]. Both siRNAs and miRNAs are generated by Dicer (DCR), a multidomain enzyme of the RNase III family. DCR cuts long, double-stranded RNA (dsRNA) into siRNAs and chops short precursor miRNAs with imperfect stem-loop structure into miRNAs.

The nascent siRNAs and miRNAs are double-stranded duplexes. These duplexes need to be unwound before they can be assembled into an RNA-induced silencing complex (RISC). By comparing the thermodynamic stabilities at the two ends, siRNAs can be divided into two classes: symmetric siRNAs and asymmetric siRNAs. A symmetric siRNA has two equally stable ends; thus, both strands of the siRNA are assembled into the RISC with equivalent efficiency [5]. By contrast, an asymmetric siRNA has one end that is less stable than the other. Because it is easier to unwind siRNA from the less stable end, one strand of the siRNA is preferentially incorporated into the RISC complex in a process referred to as the ‘asymmetric assembly of RISCs’ [5,6]. Intriguingly, most miRNAs are highly

asymmetric, ensuring efficient asymmetric assembly of a miRISC in cells [5,6].

In vitro and *in vivo* biochemical studies have shown that a siRISC can function as a miRISC to repress translation of the target mRNA; similarly, a miRISC can function as a siRISC to cleave the target mRNA. This functional interchangeability between a siRISC and a miRISC argues that siRISCs and miRISCs are highly similar, if not identical [7–10]. Much evidence suggests, however, that siRISCs and miRISCs are distinct types of complex.

First, the biogenesis, maturation and subsequent assembly of siRNAs and miRNAs into silencing complexes are different [1], which can result in RISCs with distinct functions. Second, Argonaute (AGO) proteins, which are principal components of RISCs, are encoded by a multi-gene family and can be divided into functionally distinct subgroups [11–15]. These functionally different AGOs endow their corresponding RISCs with distinct functions. Third, the complementarity between the small RNAs and their target mRNAs has been proposed to affect the functional mode of RISCs in terms of the regulation of mRNA stability and translation. Nevertheless, RISCs containing small RNAs that are extensively complementary to their target mRNAs do not always specify efficient cleavage of the targets as was previously predicted: some RISCs direct efficient target cleavage [7,8,16,17], whereas others do not [15] (G. Tang *et al.*, unpublished). Fourth, RISCs vary markedly in size, from the smallest ‘core RISC’ of ~160 kDa to the largest ‘holo RISC’ of 80 Svedberg (80S) [18–21].

Last, siRNA and miRNA programmed RISCs have distinct targeting functions in cells. Many endogenous miRNAs and their RISCs are genetically programmed to regulate gene expression and thus are important for the growth and development of an organism [22]. By contrast, siRNAs are produced from dsRNAs that are often synthesized *in vitro* or *in vivo* from viruses or repetitive sequences introduced by genetic engineering. In addition, dsRNA can be produced from endogenously activated transposons. Thus, siRNAs have been proposed to function in: (i) antiviral defense (despite the fact that viruses develop counter-defense strategies as well [23,24]), (ii) silencing mRNAs that are overproduced or translationally aborted, and (iii) guarding the genome from disruption by transposons [4,25,26].

In this review, I focus on RISC assembly by discussing the initial synthesis of siRNA and miRNA by DCRs, the formation of a RISC loading complex (RLC), the transition from an RLC to an active RISC, and target recognition and

Corresponding author: Tang, G. (guiliang.tang@umassmed.edu).

Available online 12 January 2005

Glossary

Argonaute (AGO): A large protein family that constitutes key components of RISCs. AGO proteins are characterized by two unique domains, PAZ and PIWI, whose functions are not fully understood. Current evidence suggests that the PAZ domain binds the 3'-end two-nucleotide overhangs of the siRNA duplex, whereas the PIWI domain of some AGO proteins confers slicer activity. PAZ and PIWI domains are both essential to guide the interaction between the siRNA and the target mRNA for cleavage or translational repression. Distinct AGO members have distinct functions. For example, human AGO2 programs RISCs to cleave the mRNA target, whereas AGO1 and AGO3 do not.

Cleaving RISC: A cleaving or cleavage-competent RISC that contains an endonuclease or slicer activity that binds a target for cleavage. The endonuclease or slicer identity has not been clearly identified in most organisms. Much evidence suggests that some types of AGO protein or their tightly associated identities are candidates for such an endonuclease. A cleaving RISC can direct a specific target mRNA for cleavage or translational repression, depending on the base-pairing conditions between the small RNA and the mRNA target.

Core RISC: The smallest RNA-protein complex that can direct a target mRNA for cleavage or translational repression. The RISCs of <200 kDa identified in humans and *Drosophila* might represent the core RISC. AGO proteins are associated with the core RISC.

DCR-related DExH-box RNA helicase (DRH): A protein, first identified in *C. elegans*, that has most homology to the helicase domain of DCR. There are two DRHs: DRH-1 and DRH-2. Both interact with the dsRNA-binding protein RDE-4 and are required for RNAi in *C. elegans*. Neither seems to be required for miRNA production.

Dicer (DCR): A multidomain enzyme of the RNase III family that was first identified in *Drosophila*. DCR cleaves dsRNA or stem-loop structured RNA precursors into small RNAs (~21–25 nucleotides), termed siRNAs and miRNAs. DCR is widely conserved in diverse organisms.

Drosha: A nuclear RNase III enzyme, discovered first in humans and subsequently in *Drosophila* and *C. elegans* that is implicated in initiation of the miRNA pathway.

Exportin-5: A nuclear transmembrane protein that transports precursor miRNA from the nucleus to the cytosol.

Holo RISC: The largest RNA-protein complex (80S) with RISC activity that has been found in *Drosophila*. The holo RISC associates with all possible RISC components, RLC components and proteins from other pathways. The existence of the holo RISC indicates that RISC assembly is an inseparable and sequential process and that the RISC-centered RNAi and miRNA pathways closely interact with many other pathways, possibly to control the growth and development of an organism.

Microprocessor: A nuclear complex composed of Drosha and Pasha that functions in miRNA biogenesis from the primary miRNA to the precursor miRNA.

MicroRNA (miRNA): A type of non-coding small RNA (~21–23 nucleotides) produced by DCR from a stem-loop structured RNA precursor. miRNAs are widely

expressed in animal and plant cells as RNA-protein complexes, termed miRISCs, and have been implicated in the control of development because they target specific gene transcripts for destruction or translational suppression.

Non-cleaving RISC: A RISC that lacks endonuclease activity and thus cannot target mRNA for cleavage but only for translational repression.

Pasha: A partner of Drosha in the nucleus. Pasha is a dsRNA-binding protein similar to RDE-4 and R2D2. Pasha interacts with Drosha to initiate miRNA biogenesis. In humans, Pasha is also known as DGCR8.

R2D2: A dsRNA-binding protein that has two dsRNA-binding domains and interacts with DCR2 for stability in *Drosophila*. R2D2 is thought to function as both a bridge and a sensor for functional RISC assembly.

RDE-1: An AGO protein discovered in *C. elegans*. Without RDE-1, RNAi ability is aborted (RNAi defective). RDE-1 has been proposed to interact with DCR-1, RDE-4 and DRH-1, probably for the formation of a stable protein complex that produces robust siRNAs and directs functional RISC assembly. RDE-1 does not seem to be required for miRNA production and function.

RDE-4: A dsRNA-binding protein discovered in *C. elegans*. Without RDE-4, RNAi ability is aborted (RNAi defective). RDE-4 is the homolog of R2D2. Unlike R2D2, RDE-4 has been proposed to function as the factor that presents dsRNA to DCR-1 for dicing, rather than the factor that hands siRNA to the RISC. RDE-4 does not seem to be required for miRNA production and function.

RISC loading complex (RLC): A complex that initiates formation of a RISC. The RLC sets a small RNA duplex in proper orientation for subsequent RISC assembly. Currently, siRISC-loading complexes (siRLCs) have been studied most extensively in *Drosophila*. It has been suggested that *Drosophila* siRLCs contain a DCR2-R2D2 heterodimer and the siRNA duplex; the R2D2 moiety is the asymmetric sensor that sets the siRNA orientation for RISC assembly. miRISC-loading complexes (miRLCs) have not been investigated because the biogenesis of a miRNA is more complex and an *in vitro* system for studying miRLCs has not been established.

RNA-induced initiation of transcriptional gene silencing (RITS): A complex that directs chromatin remodeling. The RITS complex also contains DCR-generated siRNA and AGO protein, and functions in heterochromatic silencing by binding to heterochromatic loci.

RNA-induced silencing complex (RISC): An RNA-protein complex that targets its perfectly or partially complementary mRNA for cleavage or translational repression. siRNA programs a siRISC and miRNA programs a miRISC. RISCs (both siRISC and miRISC) can be divided into two types: cleaving and non-cleaving. Current evidence suggests that the type of AGO protein, an essential RISC component, determines whether a RISC is cleaving or non-cleaving.

Slicer: Another term for the endonuclease located on a cleaving RISC.

Small interfering RNA (siRNA): A type of small RNA (~21–25 nucleotides) produced by DCR, a double-stranded RNA-specific enzyme of the RNase III family. The siRNA is the key component of siRISCs and triggers the silencing of its complementary mRNA.

Svedberg: Unit of sedimentation rate for measuring a particle size in a centrifuge.

interaction by the RISC. Finally, I discuss how two general types of RISC can explain the complexity of their functions in cleavage and translational repression of the target mRNA.

Initiation of the RNAi and miRNA pathways: role of DCRs

It is generally accepted that the RNA interference (RNAi) and miRNA pathways are initiated by DCR members of the RNase III enzyme family. DCR was initially identified in *Drosophila* [27] and has been subsequently found in diverse eukaryotic organisms including humans, plants and fungi (Table 1). Some organisms have a single homolog of DCR [28–35], whereas others have more [8,36]. In species with several DCRs, different homologs are responsible for producing siRNAs and miRNAs with similar or different sizes [8,21,37,38].

In *Drosophila* there are two DCR homologs, DCR1 and DCR2 (Table 1). DCR1 is responsible for miRNA production, whereas DCR2 produces siRNAs [38–40]. The fly R2D2, so named because it has two dsRNA-binding domains and tightly interacts with DCR2, has a chief role in bridging the initiation step (siRNA production) and the effector step (siRNA loading onto the effector RISC complex) of RNAi by presenting siRNA to the RLC [40,41] (Figure 1a). R2D2 and DCR2 form a stable heterodimer,

and the absence of one makes the other unstable [40]. Knocking out either DCR2 or R2D2 leads to loss of RNAi [40]. Intriguingly, whereas the *dcr2* mutant shows no obvious change in fly development, the *r2d2* mutant produces a female-sterile phenotype (Q. Liu and D. Smith, personal communication), showing that R2D2 is essential for fly development and suggesting that it has functions that are independent of DCR2.

The only DCR in *C. elegans*, DCR-1 (Table 1), is required for both the RNAi and the miRNA pathways. Genetic screening for RNAi-deficient worms has led to the identification of numerous RNAi-deficient (*rde*) genes. Biochemical studies have suggested that the dsRNA-binding protein RDE-4, an R2D2 homolog, interacts with both an AGO protein, RDE-1, and a DCR-related DExH-box RNA helicase, DRH-1. These interacting proteins work together with DCR-1 and initiate RNAi in worms [26,34] (Figure 1a). Although DCR-1 is essential for miRNA production in *C. elegans*, its interacting proteins (RDE-1, RDE-4 and DRH-1) do not seem to be involved in initiating the miRNA pathway because knocking out the genes encoding these proteins leads to no obvious abnormalities in development [34].

In contrast to the RNAi pathway, the miRNA pathway is initiated in the nucleus [42,43] (Figure 1b). In *C. elegans*,

Table 1. Principal protein components of the RLC and RISC and their putative functions

Organism	Component	Putative functions	Refs
<i>C. elegans</i>	DCR-1	Produces siRNA and miRNA	[34,75]
	Drosha	RNase III enzyme with a predominant nuclear location; interacts with Pasha to initiate the miRNA pathway	[44]
	RDE-1	A major AGO component of the RISC involved in target mRNA cleavage; interacts with DCR-1 and RDE-4 and seems to be involved in siRNA production, probably by keeping DCR-1 and RDE-4 in proper tertiary structure	[34]
	RDE-4	dsRNA-binding protein proposed to channel dsRNA to DCR1; interacts with RDE-1	[34]
	Pasha	dsRNA-binding protein (RDE-4 and R2D2 homolog) that interacts with Drosha to initiate the miRNA pathway in the nucleus	[44]
	DRH-1	RNA helicase that interacts with RDE-4 and DCR-1; is probably involved in dsRNA handling and siRNA unwinding	[34]
<i>Drosophila</i>	DRH-2	Similar to DRH-1	[34]
	DCR1	Produces miRNAs and probably interacts with AGO1	[21,27,38,40,76]
	DCR2	Produces siRNAs, interacts with R2D2 and probably interacts with AGO2	[21,38,40]
	Drosha	See entry for <i>C. elegans</i> Drosha	[44]
	R2D2	dsRNA-binding protein that binds and stabilizes DCR2, senses and binds siRNA duplexes, and probably bridges RLC to RISC	[21,40]
	Pasha	See entry for <i>C. elegans</i> Pasha	[44]
Human	AGO1	Major AGO component in miRISCs that probably interacts with DCR1; a potential weak slicer that forms a weak cleaving RISC	[39]
	AGO2	Major AGO component in siRISCs that probably interacts with DCR2; a potential slicer that forms a cleaving RISC	[39,59]
	DCR1	Produces siRNAs and miRNAs	[30,32,76]
	Drosha	See entry for <i>C. elegans</i> Drosha	[43,44]
	Pasha	See entry for <i>C. elegans</i> Pasha; also named DGCR8	[46,77]
	AGO1	An AGO component that forms a non-cleaving RISC and probably interacts with DCR1	[13]
<i>Arabidopsis</i>	AGO2	Major AGO component in siRISCs or miRISCs that probably interacts with DCR1 and functions as a slicer to form a cleaving RISC	[13,15]
	AGO3	An AGO component that forms a non-cleaving RISC and probably interacts with DCR1	[13]
	DCL1	DCR-like 1 enzyme that produces miRNAs that target endogenous mRNAs for regulation; might interact with HYL1 and HEN1 to initiate the plant miRNA pathway	[37,49]
	DCL2	DCR-like 2 enzyme; produces siRNAs that target viral RNAs	[37]
	DCL3	DCR-like 3 enzyme; produces siRNAs that direct the formation of heterochromatin	[37]
	AGO1	Major AGO component in siRISCs; a potential slicer on the cleaving RISC and component of another kind of RNA-protein complex that directs transgene methylation; probably interacts with DCL1	[62,63]
<i>S. pombe</i>	HYL1	A nuclear dsRNA-binding protein implicated in the biogenesis of plant miRNAs	[49]
	HEN1	A novel protein implicated in the biogenesis of plant miRNAs	[50]
	DCR1	Produces siRNAs and miRNAs	[28,53]
	AGO1	Major AGO component in siRISCs; a potential slicer on the cleaving RISC and a component of another kind of RITS complex that directs the formation of heterochromatin; probably interacts with DCR1	[28,53]

Drosophila and humans, an RNase III enzyme termed Drosha interacts with Pasha (a partner of Drosha with a dsRNA-binding domain similar to RDE-4 and R2D2) to form a nuclear protein complex of ~500–650 kDa termed Microprocessor. Drosha cuts the long primary miRNA transcripts into short (~70 nucleotides or longer) miRNA precursors [44–46]. These precursor miRNAs are then exported from the nucleus to the cytoplasm by Exportin-5 for maturation by cytosolic DCR [47,48]. Notably, although Pasha and DCR-1 in worms seem to be dominant in different cellular compartments, knocking down the expression of either one leads to an accumulation of primary miRNA transcripts. This suggests that there is close communication between the nuclear Microprocessor complex and cytosolic DCR-1.

Drosha and Pasha, the key components of Microprocessor, are conserved in *C. elegans*, *Drosophila* and humans but not in plants, which suggests that miRNA biogenesis diverges in different types of organism. In plants, three factors have been implicated to have a role in the biogenesis of miRNAs: DCL1, a DCR-like protein; HEN1, an uncharacterized protein; and HYL1, a nuclear dsRNA-binding protein [49,50] (Table 1). Knockout of the

genes encoding these proteins leads to a reduction in the mature miRNAs in *Arabidopsis* and an accumulation of the corresponding target mRNAs [49–51]. DCL1, HEN1 and HYL1 might interact with each other to initiate the plant miRNA pathway [49,50].

siRNA- and miRNA-protein interactions: the RLC

The DCR-cleaved siRNAs and miRNAs are initially double stranded. The transition from double-stranded to single-stranded RNAs during RISC assembly is achieved via RNA-protein and protein-protein interactions. The RLC is the initial RNA-protein complex formed in cells after the production of small RNAs. The small RNAs in the RLC are probably double stranded and ready to be unwound for functional RISC assembly. The RLC has been characterized to some extent in *Drosophila* and *C. elegans*, but its exact composition is unknown.

In *Drosophila*, formation of the RLC is initiated via interactions between the siRNA duplex and DCR2–R2D2 heterodimers [40]. A UV cross-linking study has shown that R2D2 is important in sensing the asymmetry of a siRNA duplex and selects which strand of the siRNA duplex to incorporate into the RISC (while the other




Figure 1. Models of RISC assembly: from siRNA and miRNA generation to siRISC and miRISC formation. **(a)** siRNA is generated by a complex of DCR2 (green) and R2D2 (red) in *Drosophila*, or DCR-1 (green), RDE-1 (light blue), RDE-4 (red) and DRH-1 (dark green) in *C. elegans*, and remains within the complex. In *Drosophila*, the siRNA–DCR2–R2D2 complex is thought to interact with RNA helicase, AGO protein and other RISC components to form an RLC-like complex that is rapidly converted into active RISC. In *C. elegans*, the DCR-1–RDE-1–RDE-4–DRH-1 complex seems to function in both siRNA production and subsequent assembly of the RISC. **(b)** By contrast, miRISC assembly is initiated in the nucleus, where the gene encoding the miRNA is transcribed into a primary miRNA (Pri-miRNA) transcript. The primary miRNA is processed into a miRNA precursor (Pre-miRNA) by Drosha (green) and Pasha (red) in the Microprocessor complex, and is then exported into cytoplasm by Exportin-5 (purple) for maturation into miRNA by a cytosolic DCR (DCRx; green). miRNA in the miRNA–DCR complex is subsequently assembled into the miRISC. Panel (b) is modified from Ref. [74].

strand is destroyed) [41]. This finding indicates that R2D2 not only is the DCR2 stabilizer, as mentioned above, but also is involved in functional assembly of the RISC [40,41]. How R2D2 functions in the RLC and guides asymmetric assembly of the RISC is poorly understood. R2D2 tends to bind the more stable end of the siRNA duplex and enables the siRNA duplex to unwind from the less stable end, leading to asymmetric RISC assembly [41].

Whether R2D2 or an R2D2-like sensor is required for miRISC assembly has not been fully determined. R2D2 is required for the normal development of a female fly, suggesting that it might be involved in miRNA biogenesis (Q. Liu and D. Smith, personal communication). Most mature miRNAs are detected in cells in a single-stranded form, and the hypothetical miRNA antisense strand (miRNA*) is often undetectable [50,51]. The hypothetical miRNA duplexes produced from precursor miRNA are often characterized by unequal base-pairing stability at the small duplex ends. These findings suggest that endogenous miRNAs might also use an asymmetric RISC assembly mechanism [5,6].

The mechanism of asymmetric miRISC assembly is unknown and might differ from that of asymmetric siRISC assembly. First, miRNA maturation requires two steps: the production of precursor miRNA from the primary miRNA transcript in the nucleus by Microprocessor (the Drosha–Pasha complex), followed by the maturation of miRNA in the cytosol by a specific DCR (Figure 1b). It is not known how cytosolic DCR recognizes the precursor miRNA that is transported from the nucleus or how it produces a mature miRNA in the cytosol. Second, most

miRNAs have a 5' uridine that forms a U•G wobble or mismatch base pair at the hypothetical miRNA duplex end [5,6,8]. These 5' U•G wobble or mismatch pairs might have a role in facilitating miRISC assembly.

The RLC has not been specifically studied in *C. elegans*; however, DCR-1, RDE-1, RDE-4 and DRH-1 interact with each other during RNAi initiation, which suggests that a similar complex might function as an RLC in this organism [34].

RLC structure and assembly: orientation of the small RNA duplex

In *Drosophila*, asymmetric RISC assembly requires an asymmetric RLC, which is primarily determined by the structure of the siRNA duplex [41]. The composition of the RLC has not been determined in full. The minimal components of the RLC are thought to be the siRNA duplex and the DCR2–R2D2 heterodimer [41]. It is not known what and how additional cofactors are involved in RLC assembly.

One of the key issues concerning asymmetric RLC assembly is whether the siRNA duplex produced by the DCR2–R2D2 heterodimer needs to be released and rebound to the dimer for proper orientation. Synthetic siRNA can be sensed directly and recruited by the DCR2–R2D2 heterodimer, and this hypothetical siRNA releasing step is unnecessary. UV crosslinking data have shown that the R2D2 moiety tends to bind the more stable Watson–Crick base-pairing end, whereas the DCR2 moiety tends to be crosslinked to the opposite, less stable

duplex end [41]. These results suggest that the assembly of an active RLC favors proper orientation of the siRNA.

In *Drosophila*, siRNA duplexes are produced from long dsRNA, probably in a random direction, by DCR2–R2D2. Each duplex binds one DCR2–R2D2 heterodimer, most likely via the R2D2 moiety. If the R2D2 happens to bind the more stable base-pairing duplex end, thereby positioning the DCR2 moiety at the opposite, less stable end, a highly asymmetric RLC will form [41]. How the siRNA duplex is dissociated from and rebound to DCR2–R2D2 in its proper orientation for subsequent active assembly of the RLC is unknown.

The transition from RLC to RISC: RNA helicase and AGO proteins

Handing siRNAs from the RLC to the RISC requires an AGO protein and possibly also an RNA helicase (Table 1). Even though *Drosophila* DCR2 has a DExH helicase domain, it does not unwind siRNA duplexes *in vitro* [41], which suggests that a specific RNA helicase or other factors might be required during the transition from RLC to RISC. Such a hypothetical RNA helicase is thought to unwind the siRNA duplex, after which the RISC is formed by the recruitment of an AGO protein as the core component. Indeed, purified AGO proteins from transfected cells are associated with single-stranded siRNAs or miRNAs [12,13,15,18,19], which suggests that the siRNAs or miRNAs are unwound before their stable association with AGO proteins. The helicase responsible for this unwinding has not been identified. Alternatively, other types of protein that lower the base-pairing energy might exist. These proteins might bind small RNA duplexes and help to recruit AGO proteins to one or the other single-stranded siRNA without the need of any helicase.

A common feature of RISCs from diverse organisms is that the AGO proteins are always associated with RISCs. AGO proteins constitute a large gene family in diverse organisms [11]. Most eukaryotic organisms have several AGO homologs: human has eight [52], *Arabidopsis* has ten [11], *Drosophila* has five [11] and *C. elegans* has twenty-seven [53]; however, fission yeast has only one AGO homolog [28,29,53]. It has been proposed that different resident AGO proteins might specify distinct RISC functions. This possibility is clearly demonstrated in *Drosophila*, in which AGO2 cofractionates and functionally associates with DCR2, whereas AGO1 cofractionates and functionally associates with DCR1 [39,54,55]. These results correlate with the fact that DCR1 is implicated in the maturation of miRNAs, whereas DCR2 has been shown to initiate RNAi in *Drosophila* [38,39] (Table 1). It seems that the fly AGO proteins functionally interact with different DCRs for the downstream assembly of distinct RISCs.

In contrast to *Drosophila*, humans and *C. elegans* contain only one DCR that initiates the assembly of both siRISCs and miRISCs. Nevertheless, different human AGO proteins (AGO1–AGO3) can be recruited by siRNA or miRNA to form RISCs, although only AGO2-associated RISC can direct the target mRNA for cleavage [13,15]. How one DCR transfers the small RNAs to different

AGO proteins for the assembly of distinct RISCs remains unknown.

RISC assembly from RLC: a sequential process

Recent biochemical studies have shown that RISCs vary markedly in size from complexes as small as 160 kDa in humans to ones as large as 80S in *Drosophila* [12,18–21,56]. *In vitro* RISC assembly in *Drosophila* indicates that assembly starts with the formation of an RLC that contains DCR2 and R2D2, and ends with a large protein complex the size of 80S ribosome [21]. Furthermore, a series of intermediate complexes has been observed during the transition from RLC to RISC. Notably, the RLC, the intermediates and the 80S active RISC all contain DCR2 and R2D2 [21]. The complexes that form during the transition are dependent on ATP and can be chased quantitatively on a native gel, which suggests that RISC assembly is a sequential process and that the active core RISC might be tightly associated with the RLC [21,56]. Whether such a phenomenon can be observed in other organisms by a similar experimental approach has not been tested.

Why do the RISCs seem to be so large and heterogeneous? One explanation is that these complexes might associate with ribosomes [55,57,58]. Indeed, active RISC cofractionates with ribosomes on a gel filtration column [21]. In *Drosophila*, the RISCs could be heterogeneous if the RLC and the RISC can interact with each other. For example, R2D2 in an RLC might interact with AGO proteins on a RISC. Such an interaction occurs in *C. elegans*, in which the R2D2-like protein RDE-4 physically interacts with the AGO homolog RDE-1 [34]. Intriguingly, *Drosophila* DCR1, which is implicated in the production of miRNA, also seems to associate with siRNA-programmed RLCs and RISCs [21]. This observation suggests that *Drosophila* DCR1 might share structural regions with DCR2 that enable it to interact with these complexes.

RISC functions: target mRNA cleavage and the ‘slicer’ activity

In RNAi, cleavage of an RNA target by a RISC is defined mainly by the endonuclease or ‘slicer’ activity of the RISC. Much recent evidence has demonstrated that slicer is a subtype of the AGO protein family [13,15,59]. AGO proteins have been previously shown to be in tight association with siRNAs and miRNAs by affinity purification studies using either tagged AGO protein or tagged siRNA [13,15,18,19]. Notably, among the tagged AGO proteins that have been expressed in human cells, only mammalian AGO2 has been found to program a cleavage-competent RISC *in vivo* [13,15]. Moreover, purified tagged human AGO2, together with 5′ phosphorylated single-stranded siRNA, can be forced into a core RISC-like complex that can direct target mRNA for cleavage *in vitro* [13]. This experiment indicates that human AGO2 has slicer activity. This finding is not limited to mammals. In *Drosophila*, purification of the RISC coupled with protein mass analysis has also shown that *Drosophila* AGO2 is the only protein associated with an active RISC activity [59].

Slicer activity has been mapped to the PIWI domain in AGO proteins through mutagenesis [13], a finding that is supported by the crystal structure study of an

archaebacterium protein, *Pyrococcus furiosus* AGO (Pf-AGO), and bioinformatic data [13,14,59]. The PIWI domains of Pf-AGO and *Drosophila* AGO2 are similar to RNase H or endonuclease V in secondary structure [14,59] and share conserved amino acid residues in the active site. These findings suggest that the PIWI domain of these AGO proteins confers the endonuclease activity of slicer. Several biochemical and crystal structure studies suggest that the PAZ domain of AGO proteins binds the 3' end of the single-stranded siRNA [60,61] that interacts with the target mRNA. The crystal structure of Pf-AGO suggests that the PAZ and PIWI domains define a substrate-binding groove that supports interactions between the siRNA and the target mRNA and subsequent cleavage of the target mRNA [14].

The possibility of reconstituting *in vitro* core RISC activity that can cleave a mRNA target provides a practical approach to studying the structure and function of AGO proteins in terms of slicer activity. Why do some AGOs have slicer activity and others not? Mutagenesis and bioinformatics analyses of the PIWI domain of human AGO2 and *Drosophila* AGO2 have identified key amino acid residues that are essential for slicer activity [13]. A loss of these residues during the evolution of AGO protein family might be the reason for the lack of slicer activity in some AGO proteins.

Most eukaryotes have conserved RNAi and miRNA pathways and also contain several AGO proteins. These organisms are expected to have at least one member of the AGO family with slicer activity for RNAi to function.

In *Drosophila*, both AGO1 and AGO2 are good candidates for this activity [39,59]. In *Schizosaccharomyces pombe*, only one AGO1 protein, which functions in both the RNAi and the chromatin-remodeling pathways, has been identified [28,53]. In *Arabidopsis*, AGO1 is a good candidate for slicer activity [62,63]. Experiments should be designed to test whether these candidate 'slicers', like human AGO2, can be reconstituted into core RISCs that are competent for target mRNA cleavage *in vitro*. More extensive mutagenesis and domain-swapping experiments on the PIWI domain of different AGO family members will be essential to expand our understanding of the slicer activity.

Two types of RISC: multiple functions

RISC assembly is most complex in the RNAi and miRNA pathways. It involves small RNA producers (DCRs), small RNA duplex structures and RLC assemblies, and it requires the unwinding of a symmetric–asymmetric RNA duplex and the recruitment of distinct AGO proteins. Different AGO proteins on RISCs have distinct functions that are most probably determined by the PIWI domain of the AGO protein. Moreover, some PIWI domains confer slicer activity, whereas others do not. On the basis of the type of AGO protein that is recruited to a RISC, these complexes can be tentatively divided into two general types: a cleaving RISC and a non-cleaving RISC (Figure 2).

A cleaving RISC has dual functions that can direct the target mRNA either for cleavage or for translational repression, depending on the base-pairing features between the small RNA and the target mRNA [7,8]. If




Figure 2. Model of RISC formation and function. Distinct subtypes of Argonaute protein (AGO_x and AGO_y) are proposed to interact with siRNAs or miRNAs to form two types of RISC: a cleaving and a non-cleaving RISC. A cleaving RISC can direct the target mRNA for either cleavage or translational repression, depending on the rate of target cleavage. When the small RNA (siRNA or miRNA) is extensively complementary to its target mRNA, most cleaving RISCs direct cleavage of the target mRNA at a high rate and efficiently destroy it. These cleaving RISCs can become non-cleaving if the base-pairing conditions between the small RNA and the target mRNA are not met. An extensively complementary cleaving RISC can also result in translational repression when the target cleave rate is below a certain threshold. By contrast, when bound to the target mRNA a non-cleaving RISC can direct the target mRNA only for translational repression, regardless of complementarity between the small RNA and the target.

the base-pairing complementarity between the siRNA or miRNA and its mRNA target is less extensive, the target mRNA might be physically unreachable by the active center of the endonuclease (slicer) in the cleaving RISC, because of the distorted helix that forms between the siRNA or miRNA and the target [7,64,65]. This will result in translational repression instead of efficient cleavage of the target mRNA. By contrast, non-cleaving RISCs lack endonuclease (slicer) activity in the PIWI domain of AGO proteins and can direct the target mRNA only for translational repression (Figure 2).

In addition to these two types of RISC, other kinds of RISC-like identity, such as the RNA-induced initiation of transcriptional gene silencing (RITS) complex that directs chromatin remodeling, also exist [28,66]. The RITS complex also contains DCR-generated siRNA and AGO protein, and functions in heterochromatic silencing by binding to heterochromatic loci [67]. It is not yet known whether the AGO protein on the RITS complex confers the slicer activity.

Both cleaving and non-cleaving RISCs need to recognize and to bind their targets to function in cleavage or translational regulation. Several observations indicate that target recognition and binding by a RISC are determined mainly by the base-pairing between the 5' portion of the siRNA or miRNA and its target mRNA. For example, a siRISC- or miRISC-directed target is cleaved between positions 10 and 11 from the 5' end of a siRNA or miRNA guide strand, regardless of the size (e.g. 21–25 nt) of the siRNA or miRNA [7,68]. This suggests that the 5' end (~9 nt) of a siRNA or miRNA is crucial for target recognition, binding and cleavage. In addition, bioinformatics analysis of miRNA target regions and *in vitro* target cleavage analysis of the targets with mutations in the siRNA- or miRNA-binding regions also strongly support the importance of base-pairing between the 5' portion of the siRNA or miRNA and its target [16,64,69].

Target recognition and binding by RISC do not necessarily result in target cleavage. Several factors determine whether a target gets cleaved or not. First, only cleaving RISCs have the potential to direct the targets for cleavage; a non-cleaving RISC can direct a target mRNA only for translational repression. Second, the base-pairing condition between the small RNA and its target determines whether the target mRNA is cleaved or not; in other words, a cleaving RISC might not cleave its target if the base-pairing requirement is not met. This idea is strongly supported by the fact that miRNAs and siRNAs are functionally interchangeable, depending on complementarity between the small RNAs and their targets [7,9,10]. Only extensive complementarity in specific regions results in cleavage of the target mRNA; otherwise, the RISC seems to target the mRNA for translational repression [70–72].

Third, the modes of regulating target mRNA (cleavage or translational repression) by siRNA or miRNA sometimes intersect with each other. In this case, both modes of regulation are involved. Which mode of regulation occurs *in vivo* depends on which mode is dominant. For example, miR172, an *Arabidopsis* miRNA, shows slow cleavage of its target *AP2* mRNA both *in vitro* (G. Tang *et al.*, unpublished) and *in vivo* (the 3' cleavage product can be cloned by a PCR-based approach) [71]. Generally,

however, *AP2* mRNA does not appear to be cleaved *in vivo*; instead, it seems to be translationally repressed by miR172 [70,71], indicating that miR172-directed *AP2* mRNA translational repression is predominant in *Arabidopsis*. It seems that some threshold of the rate of RISC-directed cleavage determines whether a target mRNA is destroyed or suppressed against translation.

The sequence context of a small RNA might also have a role in determining whether a cleaving or non-cleaving RISC is assembled. For example, chemically synthesized *Drosophila* miR2b, although perfectly complementary to its target mRNA, does not direct apparent cleavage of the target *in vitro*, even under conditions that are favorable for a cleaving RISC (G. Tang *et al.*, unpublished). Furthermore, the characteristics of a target mRNA (e.g. abundance, turnover, structure and translatability) might also affect the efficiency of target clearance or translational repression by specific RISCs.

Interactions between the RISC and other pathways

RISCs function in the control of specific pathways primarily by negatively regulating gene expression. Sometimes, the RISC regulates its targeting pathways by interacting with proteins from these pathways. For example, immunoprecipitation and sucrose gradient fractionation of RISCs have identified not only the common components of an active holo siRISC [21], but also proteins from other pathways [54,58,73]. These RISC interacting partners include ribosome proteins from the protein translation pathway, vasa intronic gene (VIG) and tudor staphylococcal nuclease (TSN) from RNA metabolism pathways, and fragile X-related protein (FXR) and the survival of motor neurons (SMN) complex from pathways involved in disease.

This type of interaction suggests that protein translation, RNA metabolism, some disease-related processes and most probably other unknown pathways intersect with the endogenous RNAi or miRNA regulation machinery. Potentially, this kind of interaction might be an endogenous strategy in the control of RNAi- and miRNA-regulated pathways.

Concluding remarks

In summary, RISC assembly is central to the RNAi and miRNA pathways. Biochemical and genetic studies have identified many putative RISC components, RISC interacting proteins and also the structural features of the siRNAs and miRNAs. The various components of RISCs belong to families of proteins, and each family member specifies a RISC with distinct functions. The discovery of distinct AGO proteins and the identification of slicer identity on these proteins have established two general types of RISC: a cleaving and a non-cleaving RISC (Figure 2). Whereas non-cleaving RISCs can direct only translational repression on their target mRNAs, cleaving RISCs can direct their target mRNAs for either cleavage or translational repression, depending on the base-pairing conditions between the small RNA and its target (Figure 2). The sequence context of a siRNA or a miRNA might also contribute to functional assembly of the RISC.

The variation in size and heterogeneity of RISCs reflects either complexes formed during sequential RISC

assembly or interactions between the RNAi or miRNA pathways and their regulatory components. Future directions should focus on dissecting the core RISCs to elucidate how these complexes regulate target gene expression, while also exploring the holo RISCs to understand how the RNAi and miRNA pathways interact with other cellular mechanisms.

Acknowledgements

I thank Phillip D. Zamore, in particular, for his support and for critically reading the article. Thanks also to Craig Mello, Gad Galili, Eric Lai, Qinghua Liu, Jian-Kang Zhu, Shou-wei Ding, Zuoshang Xu, Nick Rhind, Benjamin Burr, Christian Matraga and Fengang Yu for constructive comments on the article; and colleagues for sharing unpublished results. G.T. is a Charles A. King Trust Research Fellow of the Medical Foundation with funding from the Charles A. King Trust and the June Rockwell Levy Foundation.


References

- Bartel, D.P. (2004) MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell* 116, 281–297
- Meister, G. and Tuschl, T. (2004) Mechanisms of gene silencing by double-stranded RNA. *Nature* 431, 343–349
- Baulcombe, D. (2004) RNA silencing in plants. *Nature* 431, 356–363
- Mello, C.C. and Conte, D., Jr. (2004) Revealing the world of RNA interference. *Nature* 431, 338–342
- Schwarz, D.S. *et al.* (2003) Asymmetry in the assembly of the RNAi enzyme complex. *Cell* 115, 199–208
- Khvorova, A. *et al.* (2003) Functional siRNAs and miRNAs exhibit strand bias. *Cell* 115, 209–216
- Hutvagner, G. and Zamore, P.D. (2002) A microRNA in a multiple-turnover RNAi enzyme complex. *Science* 297, 2056–2060
- Tang, G. *et al.* (2003) A biochemical framework for RNA silencing in plants. *Genes Dev.* 17, 49–63
- Doench, J.G. *et al.* (2003) siRNAs can function as miRNAs. *Genes Dev.* 17, 438–442
- Zeng, Y. *et al.* (2003) MicroRNAs and small interfering RNAs can inhibit mRNA expression by similar mechanisms. *Proc. Natl. Acad. Sci. U. S. A.* 100, 9779–9784
- Carmell, M.A. *et al.* (2002) The Argonaute family: tentacles that reach into RNAi, developmental control, stem cell maintenance, and tumorigenesis. *Genes Dev.* 16, 2733–2742
- Hammond, S.M. *et al.* (2001) Argonaute2, a link between genetic and biochemical analyses of RNAi. *Science* 293, 1146–1150
- Liu, J. *et al.* (2004) Argonaute2 is the catalytic engine of mammalian RNAi. *Science* 305, 1437–1441
- Song, J.J. *et al.* (2004) Crystal structure of Argonaute and its implications for RISC slicer activity. *Science* 305, 1434–1437
- Meister, G. *et al.* (2004) Human Argonaute2 mediates RNA cleavage targeted by miRNAs and siRNAs. *Mol. Cell* 15, 185–197
- Mallory, A.C. *et al.* (2004) MicroRNA control of *PHABULOSA* in leaf development: importance of pairing to the microRNA 5' region. *EMBO J.* 23, 3356–3364
- Llave, C. *et al.* (2002) Cleavage of Scarecrow-like mRNA targets directed by a class of *Arabidopsis* miRNA. *Science* 297, 2053–2056
- Martinez, J. and Tuschl, T. (2004) RISC is a 5' phosphomonoester-producing RNA endonuclease. *Genes Dev.* 18, 975–980
- Martinez, J. *et al.* (2002) Single-stranded antisense siRNAs guide target RNA cleavage in RNAi. *Cell* 110, 563–574
- Nykanen, A. *et al.* (2001) ATP requirements and small interfering RNA structure in the RNA interference pathway. *Cell* 107, 309–321
- Pham, J.W. *et al.* (2004) A Dicer-2-dependent 80s complex cleaves targeted mRNAs during RNAi in *Drosophila*. *Cell* 117, 83–94
- Rhoades, M.W. *et al.* (2002) Prediction of plant microRNA targets. *Cell* 110, 513–520
- Pfeffer, S. *et al.* (2004) Identification of virus-encoded microRNAs. *Science* 304, 734–736
- Ding, S.W. *et al.* (2004) RNA silencing: a conserved antiviral immunity of plants and animals. *Virus Res.* 102, 109–115
- Hannon, G.J. (2002) RNA interference. *Nature* 418, 244–251
- Tabara, H. *et al.* (1999) The *rde-1* gene, RNA interference, and transposon silencing in *C. elegans*. *Cell* 99, 123–132
- Bernstein, E. *et al.* (2001) Role for a bidentate ribonuclease in the initiation step of RNA interference. *Nature* 409, 363–366
- Verdel, A. *et al.* (2004) RNAi-mediated targeting of heterochromatin by the RITS complex. *Science* 303, 672–676
- Volpe, T.A. *et al.* (2002) Regulation of heterochromatic silencing and histone H3 lysine-9 methylation by RNAi. *Science* 297, 1833–1837
- Zhang, H. *et al.* (2002) Human Dicer preferentially cleaves dsRNAs at their termini without a requirement for ATP. *EMBO J.* 21, 5875–5885
- Billy, E. *et al.* (2001) Specific interference with gene expression induced by long, double-stranded RNA in mouse embryonal teratocarcinoma cell lines. *Proc. Natl. Acad. Sci. U. S. A.* 98, 14428–14433
- Provost, P. *et al.* (2002) Ribonuclease activity and RNA binding of recombinant human Dicer. *EMBO J.* 21, 5864–5874
- Provost, P. *et al.* (2002) Dicer is required for chromosome segregation and gene silencing in fission yeast cells. *Proc. Natl. Acad. Sci. U. S. A.* 99, 16648–16653
- Tabara, H. *et al.* (2002) The dsRNA binding protein RDE-4 interacts with RDE-1, DCR-1, and a DEXH-box helicase to direct RNAi in *C. elegans*. *Cell* 109, 861–871
- Zhang, H. *et al.* (2004) Single processing center models for human Dicer and bacterial RNase III. *Cell* 118, 57–68
- Schauer, S.E. *et al.* (2002) DICER-LIKE1: blind men and elephants in *Arabidopsis* development. *Trends Plant Sci.* 7, 487–491
- Xie, Z. *et al.* (2004) Genetic and functional diversification of small RNA pathways in plants. *PLoS Biol.* 2, E104
- Lee, Y.S. *et al.* (2004) Distinct roles for *Drosophila* Dicer-1 and Dicer-2 in the siRNA/miRNA silencing pathways. *Cell* 117, 69–81
- Okamura, K. *et al.* (2004) Distinct roles for Argonaute proteins in small RNA-directed RNA cleavage pathways. *Genes Dev.* 18, 1655–1666
- Liu, Q. *et al.* (2003) R2D2, a bridge between the initiation and effector steps of the *Drosophila* RNAi pathway. *Science* 301, 1921–1925
- Tomari, Y. *et al.* (2004) A protein sensor for siRNA asymmetry. *Science* 306, 1377–1380
- Lee, Y. *et al.* (2002) MicroRNA maturation: stepwise processing and subcellular localization. *EMBO J.* 21, 4663–4670
- Lee, Y. *et al.* (2003) The nuclear RNase III Drosha initiates microRNA processing. *Nature* 425, 415–419
- Denli, A.M. *et al.* (2004) Processing of primary microRNAs by the Microprocessor complex. *Nature* 432, 231–235
- Gregory, R.I. *et al.* (2004) The Microprocessor complex mediates the genesis of microRNAs. *Nature* 432, 235–240
- Han, J. *et al.* (2004) The Drosha-DGCR8 complex in primary microRNA processing. *Genes Dev.* 18, 3016–3027
- Lund, E. *et al.* (2004) Nuclear export of microRNA precursors. *Science* 303, 95–98
- Yi, R. *et al.* (2003) Exportin-5 mediates the nuclear export of pre-microRNAs and short hairpin RNAs. *Genes Dev.* 17, 3011–3016
- Vazquez, F. *et al.* (2004) The nuclear dsRNA binding protein HYL1 is required for microRNA accumulation and plant development, but not posttranscriptional transgene silencing. *Curr. Biol.* 14, 346–351
- Park, W. *et al.* (2002) CARPEL FACTORY, a Dicer homolog, and HEN1, a novel protein, act in microRNA metabolism in *Arabidopsis thaliana*. *Curr. Biol.* 12, 1484–1495
- Reinhart, B.J. *et al.* (2002) MicroRNAs in plants. *Genes Dev.* 16, 1616–1626
- Sasaki, T. *et al.* (2003) Identification of eight members of the Argonaute family in the human genome small star, filled. *Genomics* 82, 323–330
- Sigova, A. *et al.* (2004) A single Argonaute protein mediates both transcriptional and posttranscriptional silencing in *Schizosaccharomyces pombe*. *Genes Dev.* 18, 2359–2367
- Caudy, A.A. *et al.* (2002) Fragile X-related protein and VIG associate with the RNA interference machinery. *Genes Dev.* 16, 2491–2496
- Ishizuka, A. *et al.* (2002) A *Drosophila* fragile X protein interacts with components of RNAi and ribosomal proteins. *Genes Dev.* 16, 2497–2508
- Tomari, Y. *et al.* (2004) RISC assembly defects in the *Drosophila* RNAi mutant armitage. *Cell* 116, 831–841

- 57 Jin, P. *et al.* (2004) Biochemical and genetic interaction between the fragile X mental retardation protein and the microRNA pathway. *Nat. Neurosci.* 7, 113–117
- 58 Nelson, P.T. *et al.* (2004) miRNP:mRNA association in polyribosomes in a human neuronal cell line. *RNA* 10, 387–394
- 59 Rand, T.A. *et al.* (2004) Biochemical identification of Argonaute 2 as the sole protein required for RNA-induced silencing complex activity. *Proc. Natl. Acad. Sci. U. S. A.* 101, 14385–14389
- 60 Lingel, A. *et al.* (2003) Structure and nucleic-acid binding of the *Drosophila* Argonaute 2 PAZ domain. *Nature* 426, 465–469
- 61 Yan, K.S. *et al.* (2003) Structure and conserved RNA binding of the PAZ domain. *Nature* 426, 468–474
- 62 Vaucheret, H. *et al.* (2004) The action of ARGONAUTE1 in the miRNA pathway and its regulation by the miRNA pathway are crucial for plant development. *Genes Dev.* 18, 1187–1197
- 63 Bohmert, K. *et al.* (1998) AGO1 defines a novel locus of *Arabidopsis* controlling leaf development. *EMBO J.* 17, 170–180
- 64 Haley, B. and Zamore, P.D. (2004) Kinetic analysis of the RNAi enzyme complex. *Nat. Struct. Mol. Biol.* 11, 599–606
- 65 Chiu, Y.L. and Rana, T.M. (2003) siRNA function in RNAi: a chemical modification analysis. *RNA* 9, 1034–1048
- 66 Zilberman, D. *et al.* (2003) ARGONAUTE4 control of locus-specific siRNA accumulation and DNA and histone methylation. *Science* 299, 716–719
- 67 Noma, K. *et al.* (2004) RITS acts *in cis* to promote RNA interference-mediated transcriptional and post-transcriptional silencing. *Nat. Genet.* 36, 1174–1180
- 68 Elbashir, S.M. *et al.* (2001) Functional anatomy of siRNAs for mediating efficient RNAi in *Drosophila melanogaster* embryo lysate. *EMBO J.* 20, 6877–6888
- 69 Lai, E.C. (2002) Micro RNAs are complementary to 3' UTR sequence motifs that mediate negative post-transcriptional regulation. *Nat. Genet.* 30, 363–364
- 70 Chen, X. (2004) A microRNA as a translational repressor of *APETALA2* in *Arabidopsis* flower development. *Science* 303, 2022–2025
- 71 Aukerman, M.J. and Sakai, H. (2003) Regulation of flowering time and floral organ identity by a microRNA and its *APETALA2*-like target genes. *Plant Cell* 15, 2730–2741
- 72 Reinhart, B.J. *et al.* (2000) The 21-nucleotide *let-7* RNA regulates developmental timing in *Caenorhabditis elegans*. *Nature* 403, 901–906
- 73 Caudy, A.A. *et al.* (2003) A micrococcal nuclease homologue in RNAi effector complexes. *Nature* 425, 411–414
- 74 Tang, G. and Galili, G. (2004) Using RNAi to improve plant nutritional value: from mechanism to application. *Trends Biotechnol.* 22, 463–469
- 75 Grishok, A. *et al.* (2001) Genes and mechanisms related to RNA interference regulate expression of the small temporal RNAs that control *C. elegans* developmental timing. *Cell* 106, 23–34
- 76 Hutvagner, G. *et al.* (2001) A cellular function for the RNA-interference enzyme Dicer in the maturation of the *let-7* small temporal RNA. *Science* 293, 834–838
- 77 Landthaler, M. *et al.* (2004) The human DiGeorge Syndrome Critical Region Gene 8 and its *D. melanogaster* homolog are required for miRNA biogenesis. *Curr. Biol.* 14, 2162–2167


Getting animated with parasites!

Interested in the molecular cell biology of host–parasite interactions? Then take a look at the online animations produced by *Trends in Parasitology*, one of our companion *TRENDS* journals. The pictures below are snapshots from two of our collection of animations revealing the latest advances in understanding parasite life cycles. Check them out today!



Microsporidia: how can they invade other cells?

By C. Franzen [(2004) *Trends Parasitol.* 20, 10.1016/j.pt.2004.04.009]
<http://archive.bmn.com/supp/part/franzen.html>



Interaction of *Leishmania* with the host macrophage

By E. Handman and D.V.R. Bullen [(2002) *Trends Parasitol.* 18, 332–334]
<http://archive.bmn.com/supp/part/swf012.html>